GARDEN TECH®

SEVIN®

Insect Killer

Dust Ready To Use

Guaranteed Results or your money back

Kills By Contact

PROTECT YOUR GARDEN, FLOWERS & LAWN

FROM DAMAGING PESTS

People and pets can return to the treated area once the dust has settled

won't harm PLANTS OR BLOOMS†

WHEN USED AS DIRECTED

ACTIVE INGREDIENT:
Bifenthrin 0.1%
OTHER INGREDIENTS:..... 99.9%
TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Net Weight: 3 LB. (1.36 kg)

Distributed by: TechPac, LLC.
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

RT032222IAAB

Sevin is a registered trademark of Tessenderlo Kerley, Inc.
GardenTech is a registered trademark of Gulfstream Home and Garden Inc.

EPA Reg. No. 53883-494-71004
EPA Est. 67572-GA-1, 67572-GA-2, 69703-GA-1, 69703-GA-2,
Subscript used is first letter or run code on container.

For Consumer Refund:
TechPac LLC, PO Box 95437
Palatine, IL 60095-0437

If you have a medical emergency, need consultation on diagnosis, therapy and analysis, please call 1-(866) 945-5033.
1-(800) 420-9347. If you have questions relating to product use or application, please call 1-(800) 424-9300.
If you have a product spill or transportation emergency call 1-(800) 424-9300.

QUICK FACTS

PRODUCT FACTS	• Kills over 150 insects! • Simply shake to control listed insects • Read the label before applying!	**FRUITS & VEGETABLES**	• Protects over 125 listed fruits and vegetables • Tomato, Pepper (Bell & Non-Bell), Green Bean, Green Pea, Cucumbers, Broccoli, Strawberries, Raspberries, Carrots
WHERE TO USE	Fruit and Vegetable Gardens, Ornamental Plants and Shrubs, Flowers and Ground Covers, Lawns, Home Perimeter	**KILLS**	Ants*, Aphids, Armyworms, Hornworms, Cutworms, Japanese Beetles, Stink Bugs, Cucumber Beetles

*except Fire Ants, Harvester, and Pharaoh

FIRST AID

IF SWALLOWED:	• Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-420-9347 for emergency treatment information.

NOTE TO PHYSICIAN: This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and other aquatic organisms. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.
This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.

⚠ NOTICE: Cancer - www.P65Warnings.ca.gov

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.
IMPORTANT: READ ENTIRE LABEL BEFORE EACH USE
For Outdoor Residential Use Only
This product is a ready-to-use insecticide dust, which exhibits effective control against the pests listed when used according to label directions. This product is non-injurious to most plants; even so, all varieties of plants have not been fully tested. Before treating large numbers of plants of a single variety or first treating material not previously tested, treat a small area and observe for injury prior to full scale application. Wait 5-7 days after treatment to evaluate results. Plants under stress (such as heat, drought, or frost damage) may be sensitive to this product. Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and crack-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.
For most effective results, follow directions for specific use areas.
WHERE TO APPLY: Around Home Foundations, Lawns, Shrubs, Ornamental and Flower Gardens, Fruit and Vegetable Gardens.
WHEN TO APPLY: Apply early to prevent infestation. Otherwise, apply at the first signs of insect activity or damage.

RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dust has settled.
- Do not apply to pets or pet bedding/quarters.
- Do not make applications during rain. Avoid making applications when rainfall is expected.
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- Do not apply when the wind speed is greater than 15 mph.
- Not for use on plants grown for sale or commercial use, research, or commercial seed production.
- For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains; or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Do not allow the product to enter any drain during or after application.
- Do not apply or irrigate to point of runoff.

In New York State, this product may NOT be applied to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).
In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.
All outdoor applications must be limited to spot or crack-and-crevice treatment only, except for the following permitted uses:
1. Treatment to soil or vegetation around structures;
2. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
3. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Around Homes

HOW TO APPLY: To help prevent the invasion of buildings and structures by listed pests, lightly dust a band 3 to 4 inches wide around the outside foundation wall of the home, at a rate of 8 ounces per 1000 ft². Do not apply more than 4.5 pounds per 1000 ft² per year.

On Lawns

HOW TO APPLY: Apply when insects or their damage first appear and repeat at weekly intervals or as necessary. Apply 8 ounces per 1000 ft². Sprinkle a thin uniform layer of dust on grass. Do not apply more than 4.5 pounds per 1000 ft² per year.

On Ornamentals & Flowers

HOW TO APPLY: Apply when insects or their damage first appear and repeat at weekly intervals or as necessary. Apply 8 ounces per 1000 ft²; or thoroughly apply to place a thin uniform layer of dust on both upper and lower sides of leaves, stems and flowers. Do not apply more than 4.5 pounds per 1000 ft² per year.

Pest List for Lawns, Around Homes and on Ornamentals & Flowers

Adelphocoris Plant Bugs, *Amorbia* spp., Ants (except Fire Ants, Harvester, and Pharaoh), Arizona Cricket, Artichoke Plume Moth, Asian Lady Beetle, Billbugs, Blister Beetle, Boxelder bugs, Boxwood Leafminers, Carpenter Ants (Eastern), Carpenter Ants (Western), Clearwing Moth, Common Short-tailed Cricket, Dark Mealworms, Darkling Beetle, Deer ticks, Earwigs, Eastern Striped Cricket, Elm Bark Beetle, Elm Leaf Beetle, Elm leaf rollers, Fall Field Cricket, Firebrat, Flat Grain Beetles, Four-lined Plant

Pest List for Lawns, Around Homes and on Ornamentals & Flowers continued

Bug, Fuller Rose Beetle, Garden symphylan, German Cockroach, Glassy-winged sharpshooter, Green Fruit Beetle, Holly Leafminers, Hoplia beetle, House Cricket, Indies Short-tailed Cricket, Katydid, Lace Bug, Leaffooted Bug, Lesser Grain Borers, Lycaenid pod borer, Mealybugs, Mexican Bean Beetle, Millipedes, Mormon Cricket, Northern Wood Cricket, Oak Leafminers, Periodical Cicada, Pillbugs, Potato Beetle, Raspberry Crane Maggot, Raspberry Crown Borer, Raspberry Horntail, Redhumped Caterpillar, Rose Chafer, Rose Midges, Sand Field Cricket, Scarlet Plant Bugs, Scarlet-and-Green Leafhopper, Seedcorn maggot, Sharpshooter, Silverfish, Sod Webworm, Southeastern Field Cricket, Southern Wood Cricket, Sowbugs, Spotted wing drosophila, Spring Field Cricket, Spruce budworm, Spruce Needle Leafminers, Strawberry Crown moth, Tent caterpillars, Texas Field Cricket, Tropical Cricket, Variable Field Cricket, Vocal Field Cricket, Western Striped Cricket, Western Tent Caterpillar Moth

Home Garden Vegetables And Small Fruits

HOW TO APPLY: Apply when insects or their damage first appear. Apply 1 lb (16 ounces) per 500 square feet; or thoroughly apply to place a thin, uniform layer of dust on both upper and lower sides of leaves, stems and flowers. Reapply as necessary but do not apply more than 4.5 pounds of product per 1000 ft² per year. Observe the following crop-specific use directions, including the preharvest interval (PHI) which is the number of days between last application and harvest.

GARDEN CROP	INSECT PESTS	COMMENTS	PHI
Strawberries	Aphids, Armyworms, Flea Beetles, Corn Earworm (*Heliothis* spp.), Leafrollers, Lygus Bug (*Lygus* spp.), Plant Bugs, Spittlebugs, Stink Bugs, Strawberry Clipper Weevil, Strawberry Sap Beetle	Apply when pest populations reach damaging thresholds and repeat as necessary at 7-14 day intervals.	0
Caneberries: Blackberries, Bingleberries, Dewberries, Lowberries, Marionberries, Olallieberries, Youngberries, Loganberries, Raspberries	Black Vine Weevil, Leafrollers, Orange Tortrix, Root Weevils	One application may be made pre-bloom and a second application may be made post bloom.	3
Pears	Aphids, Codling Moth, Cutworms, Green Fruitworm, Leafhoppers, Leafminers, Leafrollers, Lygus Bug (*Lygus* spp.), Plant Bugs, Plum Curculio, San Jose Scale (*Crawlers*), Stink Bugs, Tarnished Plant Bugs	Apply as necessary to maintain control using a minimum of 30-day interval.	14
Succulent Peas and Beans: Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea, Pigeon pea, Broadbean (succulent), Lima bean (green), Runner bean, Green bean, Snap bean, Wax bean, Asparagus bean, Blackeyed pea, Chinese longbean, Cowpea, Moth bean, Southern pea, Yardlong bean, Jackbean, Sword bean	Alfalfa Caterpillar, Aphids, Aster Leafhopper, Bean Leaf Beetle, Beet Armyworm, Cloverworm, Corn Earworm, Corn Rootworm, Cucumber Beetles, Cutworms, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshoppers, Japanese Beetle, Leafhoppers, Loopers, Pea Leaf Weevil, Pea Weevil, Plant Bug, Sap Beetle, Southern Armyworm, Stink Bugs, Tarnished Plant Bug, Thrips, Webworms, Western Bean Cutworm, Yellowstriped Armyworm	Do not make applications less than 7 days apart.	3
Head and Stem Brassica Vegetables: Broccoli, Chinese Broccoli (Gailon, White Flowering Broccoli), Brussels Sprouts, Cauliflower, Cavalo broccolo, Kohlrabi, Cabbage, Chinese Cabbage (napa), Chinese Mustard Cabbage (gai choy)	Aphids, Armyworms, Corn Earworm, Crickets, Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles, Ground Beetles, Imported Cabbageworm, Saltmarsh Caterpillar, Leafhoppers, Loopers, Stink Bugs, Thrips, Tobacco Budworm, Wireworm	Do not make applications less than 7 days apart. Do not make more than 5 applications after bloom.	7
Cucurbits: Chayote (fruit), Chinese Waxgourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Edible Gourd (includes hyotan, cucuzza), *Luffa* spp. (includes hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or Cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and Snake melon), Pumpkin, Summer Squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter Squash (includes Buttermut squash, calabaza, hubbard squash (*C. mixta; C. pepo*) includes acorn squash, spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus* spp.).	Aphids, Armyworms, Cabbage Looper, Cutworms, Cucumber Beetles, Corn Earworm, Grasshopper, Leafhoppers, Melonworm, Pickleworm, Plant Bug, Rindworm, Squash Bugs, Squash Vine Borer, Stink Bugs, Tobacco Budworm	Do not make applications less than 7 days apart. Do not make more than 2 applications after bloom.	3
Lettuce (Head)	Aphids, Armyworms, Corn Earworm, Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Loopers, Salt Marsh Caterpillar, Stink bug, Tobacco Budworm	Do not make applications less than 7 days apart.	7
Soybeans/Edamame	Soybean Aphid	Apply foliar treatments at a minimum of 30 days intervals.	18
Fruiting vegetables (except Tomato & Tomatillo): Eggplant, Pepper (Bell & Non-Bell), Groundcherry, Pepino	Armyworms, Cabbage Looper, Colorado Potato Beetle, Corn Earworm, Cucumber Beetle, Cutworms, European Corn Borer, Flea Beetle, Leafminers, Loopers, Pepper Weevil, Plant Bug, Stink Bug, Thrips, Tomato Pinworm, Tomato Hornworm	Do not make applications less than 7 days apart.	7
Tomato & Tomatillo	Aphids, Armyworms, Bean Leaf Beetle, Cabbageworm, Cloverworm, Corn Earworm, Corn Rootworm, Cucumber Beetles, Cutworms, Diamondback Moth, European Corn Borer, Flea Beetles, Flea Hopper, Grasshopper, Japanese Beetle, Leafhoppers, Loopers, Lygus Bug (*Lygus spp.*), Melonworm, Pea Weevil, Pea Leaf Weevil, Pickleworm, Plant Bug, Rindworm, Salt Marsh Caterpillar, Sap Beetle, Seedpod Weevil, Squash Bugs, Stink bug spp., Tobacco Budworm, Tarnished Plant Bug, Thrips, Wireworms	Do not make applications less than 10 days apart. Do not apply more than 4 applications (0.32 lb active ingredient) per season.	1
Peanut	Beet Armyworm, Corn Earworm, Cutworm spp., Fall Armyworm, Grasshoppers, Green Cloverworm, Leafhoppers, Lesser Cornstalk Borer, Loopers, Rednecked Peanut Worm, Southern Armyworm, Southern Corn Rootworm, Stink bugs, Three Cornered Alfalfa Hopper, Velvetbean Caterpillar, Yellowstriped Armyworm	At a minimum of 14 days intervals.	14
Root Crops (except Sugar Beets & Garden Beets): Burdock (edible), Carrot, Celeriac, Chervil (Turnip Rooted), Chicory, Ginseng, Horseradish, Parsley (Turnip Rooted), Parsnip, Radish, Radish (Oriental), Rutabaga, Salsify, Salsify (Black), Salsify (Spanish), Skirret, Turnip	Aphids, Beet Armyworm, Celery Leaf Tier, Corn Earworm, Cross-Striped Cabbageworm, Cutworms, Diamondback Moth, European Corn Borer, Fall Armyworm, Flea Beetles, Green Cloverworm, Hornworms, Imported Cabbageworm, Loopers, Southern Armyworm, Tobacco Budworm, Velvetbean Caterpillar, Yellowstriped Armyworm	Do not make applications less than 7 days apart.	21
Leafy Petiole Vegetables: Celery, Cardoon, Chinese Celery, Celtuce, Florence Fennel, Rhubarb, Swiss Chard	Aphids, Armyworms, Cutworms, Corn Earworm, Crickets, Cucumber Beetles, Diamondback Moth, Flea Beetles, Ground Beetles, Imported Cabbageworm, Leafhoppers, Loopers, Saltmarsh Caterpillar, Stink Bugs, Thrips, Tobacco Budworm, Wireworm	Do not make applications less than 7 days apart.	7
Bushberries: Blueberry (Highbush and Lowbush), Currant Elderberry, Gooseberry, Huckleberry	Aphids, Blueberry Maggot, Fruitworms, Leafhoppers, Lecanium Scale (crawlers), Plum Curculio, Oblique Banded Leaf Roller, Red Banded Leafroller, Spanworm, Variegated Leafroller	Do not make applications less than 7 days apart.	1

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store in tightly closed container in a dry place.
CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency or 1-866-945-5033 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Read the entire Directions For Use and Conditions of Sale and Limitation of Warranty before buying and using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

WARRANTY STATEMENT

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, User agrees to hold TechPac LLC and Seller harmless for any claims relating to such factors.

6 13499 01041 4

www.gardentech.com