



GARDEN  TECH[®]
SEVIN[®]
Sulfur Dust

Dust or Spray



Kills By Contact



Guaranteed Results or your money back



Strong Protection for Your Garden

2-IN-1 DISEASE & INSECT CONTROL

won't harm PLANTS OR BLOOMS

People and pets may re-enter once the dust settles or product dries

ACTIVE INGREDIENT:	
Sulfur*	90%
OTHER INGREDIENTS:	10%
TOTAL	100%

RT1Z2S2020C
 *Contains 0.54 lb active ingredient per gallon.
 Intended for Residential Use Only

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

See back booklet for additional
 Precautionary Statements and First Aid.

Net Weight: 1.25 LB (567 g)






DISEASE CONTROL



KILLS BY CONTACT

GARDEN TECH. **SEVIN** Sulfur Dust Dust or Spray

Quick Facts

 PRODUCT FACTS	Starts killing immediately upon contact Kills chewing and sucking insects Won't harm flowers, shrubs, or blooms Use as dust or spray	 WHERE TO USE	Fruit and vegetable gardens, ornamental plants and shrubs, flowers and ground covers
 KILLS	Controls Brown Rot, Powdery Mildew, Rust, Scab, Leaf Spot, Botrytis Blight, Black Spot, Peach Canker, Shothole, and other listed diseases Kills Thrips, Mites, Soft Scales, Citrus Red Spider, Red Spider and other listed pests	 PEOPLE & PETS	People and pets may re-enter once the dust settles or product dries
 FRUITS & VEGETABLES	Can be used on Asparagus, Beans, Carrots, Onions, Celery, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale*, Melons, Peas, Peppers, Potatoes, Spinach, Tomatoes, Avocados, Blueberries, Blackberries, Raspberries, Strawberries, Cherries, Citrus, Grapes, Mangoes, Nectarines, Peaches, Plums, and other listed fruits and vegetables *Not for use in California.		

Distributed by: TechPac, L.L.C.
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 802-16-71004
EPA Est. No. 67572-GA-1, 67572-GA-2,
69703-GA-1, 69703-GA-2
Subscript used is first letter of run
code on container.



For more information visit us
at Gardentech.com

GardenTech is a registered
trademark of Gulfstream
Home & Garden, Inc.
Sevin is a registered
trademark of Tessenderlo
Kerley, Inc.


GARDEN TECH.
SEVIN[®]
Sulfur Dust Dust or Spray

FIRST AID	
Have product container or label with you when calling a poison control center or doctor, or going for treatment.	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or on clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if inhaled, absorbed through skin, or swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water. **DO NOT** contaminate water when disposing of rinsate or equipment washwater.

STORAGE AND DISPOSAL

Storage: Store product in original container in a cool, dry, locked place out of the reach of children and pets. **Pesticide Disposal & Container Handling:** Nonrefillable container. **DO NOT** reuse or refill this container. *If empty*—Place in trash or offer for recycling if available. *If partly filled*— Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Distributed by:
 TechPac, L.L.C.
 1000 Parkwood Circle, Suite 700
 Atlanta, GA 30339

EPA Reg. No. 802-16-71004
 EPA Est. No 67572-GA-1
 67572-GA-2, 69703-GA-1,
 69703-GA-2
 Subscript used is first letter
 of run code on container.

GardenTech is a registered
 trademark of Gulfstream
 Home & Garden, Inc.
 Sevin is a registered trademark
 of Tessenderlo Kerley, Inc.



DISEASE CONTROL



KILLS BY CONTACT

GARDEN TECH® **SEVIN®** Sulfur Dust Dust or Spray

Quick Facts

PRODUCT FACTS 	Starts killing immediately upon contact Kills chewing and sucking insects Won't harm flowers, shrubs, or blooms Use as dust or spray	WHERE TO USE 	Fruit and vegetable gardens, ornamental plants and shrubs, flowers and ground covers
KILLS 	Controls Brown Rot, Powdery Mildew, Rust, Scab, Leaf Spot, Botrytis Blight, Black Spot, Peach Canker, Shothole, and other listed diseases Kills Thrips, Mites, Soft Scales, Citrus Red Spider, Red Spider and other listed pests	PEOPLE & PETS 	People and pets may re-enter once the dust settles or product dries
FRUITS & VEGETABLES 	Can be used on Asparagus, Beans, Carrots, Onions, Celery, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale*, Melons, Peas, Peppers, Potatoes, Spinach, Tomatoes, Avocados, Blueberries, Blackberries, Raspberries, Strawberries, Cherries, Citrus, Grapes, Mangoes, Nectarines, Peaches, Plums, and other listed fruits and vegetables *Not for use in California.		

Distributed by: TechPac, L.L.C.
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

EPA Reg. No. 802-16-71004
EPA Est. No 67572-GA-1, 67572-GA-2,
69703-GA-1, 69703-GA-2,
Subscript used is first letter of run
code on container.



For more information visit us at
Gardentech.com

GardenTech is a registered
trademark of Gulfstream
Home & Garden, Inc.
Sevin is a registered trademark
of Tessenlerio Kerley, Inc.

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347

If you have questions relating to product use or application, please call (866) 945-5033

If you have a product spill or transportation emergency call (800) 424-9300

For Consumer Refund: TechPac, LLC, PO Box 95437 Palatine, IL 60095-0437

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Intended for residential use only.

This product is a combination fungicide and insecticide that can be used as a dust or spray. It can be used on listed flowers, ornamentals, fruits, and vegetables. It kills listed pests and controls diseases including powdery mildew, leaf spot, black spot, scab, rust, and more listed below.

HOW TO USE

Please read the entire label. Use only as directed, noting all use restrictions. Consult State Agricultural Experiment Station or the State Agricultural Extension Service for sensitive plants and additional information as the timing, number and rate of application needed may vary with local conditions.

AS A DUST: Before dusting, shake duster vigorously to mix air with the sulfur. Dust lightly. Frequent light dustings are more effective than single heavy dustings.

AS A SPRAY: Remove cap with dip tube to measure dust. Mix the required amount of Sevin® Sulfur Dust with sufficient water to make a thin, creamy paste before diluting with the balance of the required amount of water. Use about 1 gallon of diluted spray per 1000 square feet.

tsp = teaspoon tbsp = tablespoon

Fruits & Nuts

Make full coverage applications on both sides of leaves. Apply to plant foliage, stems, and flowers to leave a "barely visible" dust. As a preventive measure or for light infestations, use the lower dosage rates.

USE RESTRICTIONS: DO NOT apply Sevin® Sulfur Dust dusts or sprays when shade temperatures during the day are likely to exceed 85° - 90° F. Injury to foliage and fruit may result. **DO NOT** use within 30 days of an oil spray (60 days on citrus). **DO NOT** combine with an oil spray.

Garden Crop	Target Pests/ Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Almonds	Brown Rot, Scab, Powdery Mildew, Leaf Spot, Mites (Blister, European Red, Flat, Silver, Twospotted), Rust, Twig Blight	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 12 applications per year. Minimum retreatment interval: 30 days.
Apples, Pears	Scab, Powdery Mildew, Mites, (Blister, European red, Silver, Twospotted), Apple Cedar Rust, Black Rot, Frog Eye, Quince Rust, Sooty Blotch	Spray (2 ½ - 7 ¾ tbsp per gallon of water) when buds begin to swell and thereafter until 2 weeks after the petals fall. Treat Delicious Apples and Anjou Pears only before bloom or injury may occur.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days.
Avocados	Mites, Avocado Thrips*	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days.
Bushberries (Blueberries, Currants, Huckleberries)	Powdery Mildew	Dust or spray (1 ½ - 3 ¾ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.3 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.8 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days.
Canefruits (Blackberries, Boysenberries, Dewberries, Gooseberries, Loganberries, Raspberries)	Powdery Mildew	Dust or spray (1 ½ - 3 ¾ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.3 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.8 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days.
Strawberries	Powdery Mildew	Dust or spray (1 ½ - 2 ½ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days.

*Not for use in California.

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Cherries	Powdery Mildew, Silver Mite, Brown Rot, Cherry Leaf Spot, Rust, Brown Rot Blossom Blight, Leaf Spot, Mites, Red Spider, Scab, and Stem Blight	Dust or spray (2 ½ - 7 ¼ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods. The first three applications will also provide control of Brown Rot Blossom Blight. Weekly applications from 5 weeks before and continuing up to the day of harvest will provide control of Brown Rot of Fruit.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 10 days.
Citrus (Oranges, Tangelos, Tangerines, Grapefruit, Lemons, Limes, Kumquats, Citron-Citrus)	Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Six-Spotted Mite, Citrus Rust Mite, Mites, Scales (Soft, Red, Yellow, Purple, Black, and Citricola), Brown Rot, Rust Mite, Two-Spotted Mite, Citrus Red Spider, Thrips	Dust or spray (2 ¼ - 5 ¼ tbsp per gallon of water) when pests first appear. Avoid application during or preceding high temperatures. DO NOT apply within two months of an oil application.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 6.8 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 17 applications per year. Minimum retreatment interval: 21 days.
Dates	Banks Grass Mites	Dust or spray (5 ¼ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days.
Figs	Mites	Dust or spray (1 ½ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 1.7 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 17 applications per year. Minimum retreatment interval: 21 days.
Olives	Mites	Dust or spray (2 ½ - 7 ¼ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days.
Macadamias, Walnuts	Mites	Dust or spray (2 ½ - 5 ¼ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days.

*Not for use in California.

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Pecans	Mites	Dust or spray (1 ½ - 2 ½ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 4.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 21 applications per year. Minimum retreatment interval: 10 days.
Pomegranates	Mites	Dust or spray (¾ - 2 ½ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 3.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 17 applications per year. Minimum retreatment interval: 21 days.
Grapes	Powdery Mildew, Mites, Rust, Phomopsis	Dust or spray (½ - 2 ¾ tbsp per gallon of water) during the growing season when new shoot growth is 6 to 10 inches long. Concord and other labrusca type grapes may be injured by Sevin® Sulfur Dust.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.8 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 14 applications per year. Minimum retreatment interval: 26 days.
Mangoes	Powdery Mildew	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) when pests first appear. Repeat at bloom, at fruit set, and 3 weeks later.	DO NOT exceed 0.6 lb. of produce per 1000 sq. ft. per application. DO NOT exceed a total of 2.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 4 applications per year. Make four applications, as directed in Application Directions.
Nectarines, Peaches,	Brown Rot Blossom Blight, Coryneum Blight, Mites, Powdery Mildew, Silver Mite, Brown Rot, Cherry Leaf Spot, Rust, Scab, Leaf Spot, Peach Canker, Shothole, Red Spider	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods. The first three applications will also provide control of Brown Rot Blossom Blight. Weekly applications from 5 weeks before and continuing up to the day of harvest will provide control of Brown Rot of Fruit.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 10 days.
Papaya	Mites (except Carmine mites)	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days.
Peanuts	Leaf Spot, Powdery Mildew, Rust	Dust or spray (1 ½ - 2 tbsp per gallon of water) when disease first appears. Apply when disease threatens during rainy periods.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days.

Garden Crop	Target Pests/ Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Pistachios	Citrus Flat Mite, Mites	Dust or spray (2 ½ - 5 ¼ tbsp per gallon of water) when mites first appear.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days.
Plums, Prunes	Brown Rot Blossom Blight, Leaf Spot, Mites, Powdery Mildew, Silver Mite, Brown Rot, Cherry Leaf Spot, Rust, Scab, Shot-hole, Red Spider	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods. Weekly applications from 5 weeks before and continuing up to the day of harvest will provide control of Brown Rot of Fruit.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. See Application Directions for retreatment schedule to control Brown Rot of Fruit. For other uses, minimum retreatment interval: 10 days.
Quince	Brown Rot Blossom Blight, Powdery Mildew	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Sugar beets	Powdery Mildew, Red Spider Mites	Dust or spray (2 ½ - 5 ¼ tbsp per gallon of water) when disease first appears. For spider mites, dust or spray (2-11 tbsp/gallon) when pests first appear.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

Vegetables

Make full coverage applications on both sides of leaves. Apply to plant foliage, stems, and flowers to leave a "barely visible" dust. As a preventive measure or for light infestations, use the lower dosage rates.

USE RESTRICTIONS: DO NOT apply Sevin® Sulfur Dust dusts or sprays when shade temperatures during the day are likely to exceed 85° - 90° F. Injury to foliage and fruit may result. **DO NOT** use within 30 days of an oil spray. **DO NOT** combine with an oil spray.

Garden Crop	Target Pests/ Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Artichoke	Leaf Spot	Dust or spray (2 ½ - 5 ¼ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Asparagus	Rust, Mites	Dust or spray (2 ½ - 7 ¾ tbsp per gallon of water) when pests first appear and repeat through fern growth season.	DO NOT exceed 0.6 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Beans including Adzuki, Broad, Dry, Jackbean, Kidney, Lablab, Lima, Moth, Mung, Navy, Pink, Pinto, Rice, Runner, Snap, Tepary, Urd, Wax, Yardlong Peas including Blackeyed, Chickpea, Cowpea, Crowder, Dry, Dwarf, Edible Podded, English, Field, Garden, Green, Pigeon, Shelled, Snow, Southern, Succulent, Sugar Snap, Unshelled	Powdery Mildew, Rust, Thrips, Spider Mites, Leaf Spot, Mites	Dust or spray (¾ - 2½ tbsp per gallon of water) when pests are first noticed. For Rust control, start applications in early spring before any rust is noticed. Repeat applications at 10 day intervals until bloom.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Beets, Table	Mites, Powdery Mildew	Dust or spray (¾ - 1½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Carrots, Garlic, Onions, Shallots	Petrobia Mite, Powdery Mildew, Downy Mildew*	Dust or spray (¾ - 2½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Celery, Endive, Escarole, Swiss Chard, Salsify	Mites, Powdery mildew, Rust	Dust or spray (1 - 1½ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Cucumbers	Powdery Mildew	Dust or spray (½ - 1 tbsp per gallon of water) when pests first appear. May cause injury when used at higher rates during excessive temperatures. Use lower rate (½ tbsp) when temperatures exceed 85°F.	DO NOT exceed 0.08 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.1 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Cole Crops including Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Mites, Powdery Mildew, Septoria Leaf Spot	Dust or spray (¾ - 2½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Eggplant	Powdery Mildew	Dust or spray (1 - 1 ½ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Greens* including Collards, Kale, Mustard Greens	Mites, Powdery Mildew	Dust or spray (¾ - 2½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Horseradish*, Parsnips*, Radishes*, Rutabagas*, Turnips*	Powdery Mildew, Mites	Dust or spray (¾ - 2½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Lettuce* (Head & Leaf)	Powdery Mildew, Red Spider Mite, Rust	Dust or spray (1 ½ - 2½ tbsp per gallon of water) when at early leaf stage.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Melons including Watermelon, Cantaloupe, Muskmelon	Powdery Mildew	Dust or spray (5 ¼ - 6 ¾ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed ½ lb. product per 1000 sq. ft. per application. DO NOT exceed a total of 13 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Okra	Leaf Spot, Mites, Powdery Mildew	Dust or spray (¾ - 2½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days

*Not for use in California.

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Peppers (Anaheim, Banana, Bell, Cayenne, Chili, Ghost, Habanero, Jalapeno, Peppercorn, Poblano, Scotch Bonnet, Serrano, Sweet)	Leaf Spot, Powdery Mildew	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Peppermint Tops, Spearmint Tops	Powdery Mildew	Dust or spray (1 - 1½ tsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 15 days
Potatoes	Leaf Spot, Powdery Mildew, Two-spotted mites	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Pumpkins, Squash* (Summer & Winter)	Powdery Mildew	Dust or spray (1 ½ - 2 ½ tsp per gallon of water) when disease first appears. Avoid application during or preceding high temperatures.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Soybeans	Leaf Spot, Powdery Mildew, Rust	Dust or spray (2 ½ - 3 ¾ tsp per gallon of water) when disease first appears.	DO NOT exceed 0.3 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.8 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Spinach	Powdery Mildew, Rust	Dust or spray (1 - 1 ½ tsp per gallon of water) when pests are first noticed. May cause injury to certain varieties of spinach. Apply to a small area first to test for phytotoxicity before making a full application.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Sweet Potatoes	Leaf Spot, Mites, Powdery Mildew, Red Spider Mites	Dust or spray (1 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.1 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Tomatoes	Tomato Russet Mite, Mites, Powdery Mildew	Dust or spray (2 ½ - 5 ¼ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.4 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

Ornamental Flowers, Shrubs, & Trees

To control black spot and red spiders on roses, ornamental shrubs, and perennials (except in California). To control rust and powdery mildew on ornamental flowers, shrubs and trees, dust or spray as directed to completely cover foliage. As a preventive measure or for light infestations, use the lower dosage rates.

USE RESTRICTIONS: DO NOT apply Sevin® Sulfur Dust dusts or sprays when shade temperatures during the day are likely to exceed 85° - 90° F. Injury to foliage and fruit may result. **DO NOT** use within 30 days of an oil spray. **DO NOT** combine with an oil spray.

Follow directions given for specific plants.

Ornamental Flowers, Shrubs & Trees	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Alder, Azalea, Buttonbush, Catalpa, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden Fleece, Hibiscus, Honeysuckle, Horse chestnut, Lady's mantle, Ligustrum, Lilac, Linden, Matrimony vine, Crepe myrtle, Poplar, Rhododendron, Staghorn sumac, Silver vine, Spiraea, Sweetpea, Sunflower, Sycamore, Trumpetvine, Verbena, Willow, Witch hazel, Zinnia	Powdery Mildew	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Arborvitae, Cedar, Sweetpea	Red Spider Mites	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Aster	Powdery Mildew, Rust*	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Calendula, Cherry laurel, Columbine, English ivy, Foxglove, Petunia, Sage, Smilax	Leaf Spot	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Carnation	Red Spider Mites, Rust*	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Chrysanthemum	Powdery Mildew, Red Spider Mites, Rust*	Dust or spray (¾ - 2 ½ tsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

Ornamental Flowers, Shrubs & Trees	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Clematis	Leaf Spot, Red Spider Mites	Dust or spray ($\frac{1}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Hydrangea	Botrytis Blight, Leaf Spot, Powdery Mildew, Red Spider Mites	Dust or spray ($\frac{1}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Phlox	Leaf Spot, Powdery Mildew	Dust or spray ($\frac{1}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Hollyhocks, Snapdragons	Rust	Dust or spray (1 tablespoon per gallon of water) in early spring.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Roses	Powdery Mildew, Black Spot, Red Spider Mites, Rust, Brown Canker	Dust or spray (4 tbsp per gallon of water) in spring when young growth first appears.	DO NOT exceed 0.2 lb. of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

Filled by weight not by volume. Product may settle after packaging.

STORAGE AND DISPOSAL

Storage: Store product in original container in a cool, dry, locked place out of the reach of children and pets. **Pesticide Disposal & Container Handling:** Nonrefillable container. **DO NOT** reuse or refill this container. *If empty*—Place in trash or offer for recycling if available. *If partly filled*—Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

